

A Novel Mechanism to Generate FFLO States in Holographic Superconductors

James Alsup*

*Computer Science, Engineering and Physics Department,
The University of Michigan-Flint, Flint, MI 48502-1907, USA*

Eleftherios Papantonopoulos†

Department of Physics, National Technical University of Athens, GR-15780 Athens, Greece

George Siopsis‡

Department of Physics and Astronomy, The University of Tennessee, Knoxville, TN 37996 - 1200, USA

(Dated: October 5, 2012)

We discuss a novel mechanism to set up a gravity dual of FFLO states in strongly coupled superconductors. The gravitational theory utilizes two $U(1)$ gauge fields and a scalar field coupled to a charged AdS black hole. The first gauge field couples with the scalar sourcing a charge condensate below a critical temperature, and the second gauge field provides a coupling to spin in the boundary theory. The scalar is neutral under the second gauge field. By turning on an interaction between the Einstein tensor and the scalar, it is shown that, in the low temperature limit, an inhomogeneous solution possesses a higher critical temperature than the homogeneous case, giving rise to FFLO states.

PACS numbers: 11.25.Tq, 04.70.Bw, 71.45.Lr, 71.27.+a

The AdS/CFT correspondence, which was discovered in string theory, has opened up a broad avenue for the exploration of condensed matter systems at strong coupling. By using a holographic principle, these systems (described by gauge field theories) are mapped onto weakly coupled gravitational systems of one additional dimension, in which physical quantities can be computed. This holographic principle (gauge theory / gravity duality) has been applied to the study of conventional and unconventional superfluids and superconductors [1], Fermi liquids [2], and quantum phase transitions [3].

The high- T_c superconductors, such as cuprates and iron pnictides, are examples of unconventional superconductors which exhibit competing orders that are related to the breaking of the lattice symmetries. This breaking introduces inhomogeneities and a study of the effect of inhomogeneity of the pairing interaction in a weakly coupled BCS system [4] suggests that inhomogeneity might play a role in high- T_c superconductivity. In an effort to explain this behavior a “striped” superconductor was proposed [5]. Holographic striped superconductors were discussed in [6] where a modulated chemical potential was introduced and it was shown that below a critical temperature superconducting stripes develop. Properties of the striped superconductors and back reaction effects were studied in [7, 8]. Striped phases were also found in electrically charged RN-AdS black branes that involve neutral pseudo-scalars [9].

Inhomogeneous phases also appear when a strong external magnetic field coupled to the spins of the conduc-

tion electrons is applied to a high-field superconductors. This results to a separation of the Fermi surfaces corresponding to electrons with opposite spins (for a review see [10]). If the separation is too high, the pairing is destroyed and there is a transition from the superconducting state to the normal one (paramagnetic effect). An intriguing new state of matter at the transition point was proposed by Fulde and Ferrell [11] and Larkin and Ovchinnikov [12] (the FFLO state) but it has not been found experimentally so far. This state is characterized by a space modulated order parameter, corresponding to an electron pair having nonzero total momentum.

A way to understand the formation of the FFLO phase in a superconductor-ferromagnetic system (S/F) is to use the generalized Ginzburg-Landau expansion. In order to describe the paramagnetic effect in the presence of a strong external magnetic field, the usual $|\psi|^4$ -Ginzburg-Landau functional has to be modified with coefficients in the functional which depend also on the magnetic field. In this case, the (B, T) phase diagram exhibits a different behavior indicating that the minimum of the functional does not correspond to a uniform state, and a spatial variation of the order parameter decreases the energy of the system. To describe such a situation, it is necessary to add a higher-order derivative term in the expansion of the Ginzburg-Landau functional (for a detailed account see [13]).

There are several studies of the behavior of holographic superconductors in the presence of an external magnetic field. Non-trivial spatially dependent solutions have been found, like the droplet [14] and vortex solutions with integer winding number [15–17]. An analytic study on holographic superconductors in an external magnetic field was carried out in [18]. In a model resulting from a consistent truncation of type IIB string theory, anisotropic

* jalsup@umflint.edu

† lpapa@central.ntua.gr

‡ siopsis@tennessee.edu

solutions at low temperature were found [19], showing similarity between the phase diagrams of holographic superfluid flows and those of ordinary superconductors with imbalanced chemical potential. A holographic superconducting model with unbalanced fermi mixtures at strong coupling was discussed in [20]. The charge and spin transport properties of the model were studied, but the phase diagram did not reveal the occurrence of FFLO-like inhomogeneous superconducting phases.

In our recent work [21], we proposed a gravity dual of FFLO states in strongly coupled superconductors. The gravity sector consisted of two $U(1)$ gauge fields and a scalar field. The first gauge field had a non-zero scalar potential term which was the source of the charge condensate in the boundary theory through its coupling to the scalar field. The second $U(1)$ gauge field corresponded to an effective magnetic field acting on the spins in the boundary theory. The scalar field was neutral under the second $U(1)$ gauge field. We looked first at the behavior of the system at or above the critical temperature. The Einstein-Maxwell system was solved by a dyonic black hole with electric and magnetic charges, as in [20]. At the critical temperature, the system underwent a second-order phase transition and the black hole acquired hair. To find the critical temperature, we solved the scalar equation in the background of the dyonic black hole, and found that the system possessed inhomogeneous solutions for the scalar field, which however always gave a critical temperature lower than the one of the homogeneous solution. Therefore the homogeneous solutions was always dominant.

Next, we turned on an interaction term of the magnetic field to the scalar field of the generalized Ginzburg-Landau gradient type (in a covariant form). The scalar field equation was modified and the resulting inhomogeneous solutions gave a critical temperature which was higher than the one of the homogeneous solutions. We attributed this behavior of the system to the appearance of FFLO states. We noted that the appearance of the FFLO states was more pronounced as $T_c/\mu \rightarrow 0$, and the magnetic field of the second $U(1)$ gauge group was large.

In this letter, we propose a novel mechanism for the generation of the gravity dual of FFLO states in the low temperature limit. In our previous work we showed that, in order to generate the FFLO phase, we needed a direct coupling of the magnetic field to the scalar field. We will show that this interaction term can be effectively generated through the coupling of the Einstein tensor to the scalar field. The reason is that since the electromagnetic fields backreact on the metric, the Einstein tensor has encoded the information of these fields.

As before, the bulk theory consists of two $U(1)$ gauge fields and a scalar field. The first gauge field has a non-zero scalar potential term and the second $U(1)$ gauge field corresponds to a chemical potential (imbalance) for spin. The scalar field is neutral under the second $U(1)$ gauge field. Note, too, the second $U(1)$ is self-dual under $\vec{E} \leftrightarrow \vec{B}$ and alternatively the boundary theory can be

understood in terms of a magnetic field instead of the chosen spin chemical potential.

We turn on a new interaction between the Einstein tensor and the scalar field. Such an interaction is most often seen in scalar-tensor theories. The interest stems from the galilean symmetry of the system where the action is invariant under shifts of field derivatives by a constant vector. Thereby the higher-derivative theory has only second order equations of motion [22]. It was shown that this term acts as an effective cosmological constant and produces an early entrance into a quasi-de Sitter stage as well as a smooth exit [23]. Cosmic evolution for vanishing cosmological constant has also been investigated in [24]. The coupling has been realized in string cosmology from an effective heterotic action, up to α' corrections [25] and also in $N = 1$ four-dimensional new-minimal supergravity theories [26].

Moreover, interest away from cosmology has developed as the interaction has been used to study phase transitions for vanishing cosmological constant [27] and effects on conventional holographic superconductors employing anti-de Sitter space [28]. The presence of this term modifies the scalar field equation and the resulting inhomogeneous solutions give a critical temperature which is higher than the homogeneous solutions. We attribute this behavior to FFLO states. Note that as before, the appearance of the FFLO states is more pronounced as $T_c/\mu \rightarrow 0$ and the gauge field of the second $U(1)$ gauge group is near its maximum value.

Consider the action

$$S = \int d^4x \sqrt{-g} \left[\frac{R + 6/L^2}{16\pi G} - \frac{1}{4} F_{AB} F^{AB} - \frac{1}{4} \mathcal{F}_{AB} \mathcal{F}^{AB} \right], \quad (1)$$

where $F_{AB} = \partial_A A_B - \partial_B A_A$, $\mathcal{F}_{AB} = \partial_A \mathcal{A}_B - \partial_B \mathcal{A}_A$ are the field strengths of the $U(1)$ potentials A_A and $\mathcal{A}_A$, respectively. We set $L = 8\pi G = 1$.

The Einstein-Maxwell equations admit a solution which is a four-dimensional AdS black hole of two $U(1)$ charges,

$$ds^2 = \frac{1}{z^2} \left[-h(z) dt^2 + \frac{dz^2}{h(z)} + dx^2 + dy^2 \right], \quad (2)$$

with the horizon radius set at $z = 1$.

The two sets of Maxwell equations admit solutions of the form, respectively,

$$A_t = \mu(1 - z), \quad A_z = A_x = A_y = 0, \quad (3)$$

and

$$\mathcal{A}_t = \delta\mu(1 - z), \quad \mathcal{A}_z = \mathcal{A}_x = \mathcal{A}_y = 0, \quad (4)$$

with corresponding field strengths having non-vanishing components for electric fields in the z -direction, respectively,

$$F_{tz} = -F_{zt} = \mu, \quad \mathcal{F}_{tz} = -\mathcal{F}_{zt} = \delta\mu. \quad (5)$$

Then from the Einstein equations we obtain

$$h(z) = 1 - \left(1 + \frac{\mu^2 + \delta\mu^2}{4}\right) z^3 + \frac{\mu^2 + \delta\mu^2}{4} z^4. \quad (6)$$

The Hawking temperature is

$$T = -\frac{h'(1)}{4\pi} = \frac{3}{4\pi} \left[1 - \frac{\mu^2 + \delta\mu^2}{12}\right]. \quad (7)$$

In the limit $\mu, \delta\mu \rightarrow 0$ we recover the Schwarzschild black hole.

We now consider a scalar field ϕ , of mass m , and $U(1)^2$ charge $(q, 0)$, coupled to the Einstein tensor. The action is

$$S = \int d^4x \sqrt{-g} [(g^{AB} - \xi G^{AB}) (D_A \phi)^* D_B \phi - m^2 |\phi|^2], \quad (8)$$

where $D_A = \partial_A + iqA_A$ and G_{AB} is the Einstein tensor.

Firstly, we consider the conventional case setting $\xi = 0$. The asymptotic behavior (as $z \rightarrow 0$) of the scalar field is

$$\phi \sim z^\Delta, \quad \Delta(\Delta - 3) = m^2. \quad (9)$$

For a given mass, there are, in general, two choices of Δ ,

$$\Delta = \Delta_\pm = \frac{3}{2} \pm \sqrt{\frac{9}{4} + m^2}, \quad (10)$$

leading to two distinct physical systems.

As we lower the temperature, an instability arises and the system undergoes a second-order phase transition with the black hole developing hair. This occurs at a critical temperature T_c which is found by solving the scalar wave equation in the above background,

$$\partial_z^2 \phi + \left[\frac{h'}{h} - \frac{2}{z}\right] \partial_z \phi + \frac{1}{h} \nabla_2^2 \phi - \frac{1}{h} \left[\frac{m^2}{z^2} - q^2 \frac{A_t^2}{h}\right] \phi = 0, \quad (11)$$

with the metric function h given in (6) and the electrostatic potential A_t in (3).

Although the wave equation (11) possesses (x, y) -dependent solutions, the symmetric solution dominates and the hair that forms has no (x, y) dependence. To see this, let us introduce (x, y) -dependence and consider a static scalar field which is an eigenstate of the two-dimensional Laplacian,

$$\nabla_2^2 \phi = -\tau \phi, \quad \tau > 0. \quad (12)$$

For example, if ϕ varies sinusoidally in the x -direction, $\phi \sim e^{iQx}$, then $\tau = Q^2$ and the modulation is realized in the boundary CFT through the order parameter $\langle \mathcal{O} \rangle \sim e^{iQx}$. It is also possible for ϕ to be rotationally symmetric in the (x, y) plane, $\phi \sim J_0(\sqrt{\tau(x^2 + y^2)})$. For $\tau = 0$, we recover the homogeneous solution.

Upon factoring out the (x, y) dependence, the scalar field is represented by $\psi(z)$ and the wave equation becomes

$$\psi'' + \left[\frac{h'}{h} - \frac{2}{z}\right] \psi' - \frac{\tau}{h} \psi - \frac{1}{h} \left[\frac{m^2}{z^2} - q^2 \frac{A_t^2}{h}\right] \psi = 0. \quad (13)$$

Before we proceed with a discussion of solutions, notice that there is a scaling symmetry

$$\begin{aligned} z &\rightarrow \lambda z, \quad x \rightarrow \lambda x, \quad \tau \rightarrow \tau/\lambda^2, \\ \mu &\rightarrow \mu/\lambda, \quad \delta\mu \rightarrow \delta\mu/\lambda, \quad T \rightarrow T/\lambda, \end{aligned} \quad (14)$$

so we should only be reporting on scale-invariant quantities, such as T/μ , $\delta\mu/\mu$, τ/μ^2 , etc. It is convenient to introduce the scale-invariant parameter

$$\beta = \frac{\delta\mu}{\mu} \quad (15)$$

to describe the effect of the chemical potential imbalance.

The system is defined uniquely by specifying the parameters q and Δ . One can then vary the other parameters to study the behavior of the system. For fixed values of the scale-invariant parameters τ/μ^2 and β , we can solve the wave equation (13) and obtain $\mu = \mu_c$ as an eigenvalue. Then from (7), we deduce the critical temperature at which the second-order phase transition occurs,

$$\frac{T_c}{\mu} = \frac{T}{\mu_c} = \frac{3}{4\pi\mu_c} \left[1 - \frac{\mu_c^2(1 + \beta^2)}{12}\right]. \quad (16)$$

The maximum critical temperature is obtained for $\beta = 0$. In this case, we recover the Reissner-Nordström black hole. As we increase β , the temperature (16) decreases. For a given $\beta > 0$, the black hole is of the Reissner-Nordström form with effective chemical potential

$$\mu_{\text{eff}}^2 = \mu_c^2(1 + \beta^2). \quad (17)$$

The scalar wave equation is the same as its counterpart in a Reissner-Nordström background, but with effective charge

$$q_{\text{eff}}^2 = \frac{q^2}{1 + \beta^2}, \quad (18)$$

so that $q_{\text{eff}}\mu_{\text{eff}} = q\mu_c$.

It is known [1] that the instability [29] occurs for all values of q_{eff} , including $q_{\text{eff}} = 0$, if $\Delta \leq \Delta_*$, where $\Delta_* = \Delta_+$ for $m^2 = -\frac{3}{2}$, or explicitly,

$$\Delta_* = \frac{3 + \sqrt{3}}{2} \approx 2.366. \quad (19)$$

For $\Delta \leq \Delta_*$, β can increase indefinitely. The critical temperature has a minimum value and as $\beta \rightarrow \infty$, T_c diverges.

For $\Delta > \Delta_*$, q_{eff} has a minimum value at which the critical temperature vanishes and the black hole attains extremality. This is found by considering the limit of the near horizon region [30, 31]. One obtains

$$q_{\text{eff}} \geq q_{\text{min}} \quad , \quad q_{\text{min}}^2 = \frac{3 + 2\Delta(\Delta - 3)}{4}. \quad (20)$$

At the minimum ($T_c = 0$), $\mu_{\text{eff}}^2 = 12$, and β attains its maximum value,

$$\beta \leq \beta_{\text{max}} \quad , \quad \beta_{\text{max}}^2 = \frac{q^2}{q_{\text{min}}^2} - 1 \quad . \quad (21)$$

This limit is reminiscent of the Chandrasekhar and Clogston limit [32] in a S/F system, in which a ferromagnet at $T = 0$ cannot remain a superconductor with a uniform condensate.

In the inhomogeneous case ($\tau \neq 0$), the above argument still holds with the replacement $m^2 \rightarrow m^2 + \tau$. The effect of this modification is to increase the minimum effective charge to

$$q_{\text{min}}^2 = \frac{3 + 2\Delta(\Delta - 3) + 2\tau}{4}, \quad (22)$$

and thus decrease the maximum value of β (21). We always obtain a critical temperature which is lower than the corresponding critical temperature (for same β) in the homogeneous case ($\tau = 0$).

Notice also that $\tau \leq \tau_{\text{max}}$, where the maximum value is attained when $q_{\text{min}} = q$ (so that $\beta_{\text{max}} = 0$). We deduce from (22),

$$\tau_{\text{max}} = 2q^2 - \frac{3}{2} - \Delta(\Delta - 3). \quad (23)$$

Now let us consider the effect of coupling to the Einstein tensor by setting $\xi \neq 0$. The wave equation is modified to

$$\psi'' + \left[\frac{h'}{h} + \frac{f'_+}{f_+} - \frac{2}{z} \right] \psi' - \frac{\tau}{h} \frac{f_-}{f_+} \psi - \frac{1}{h} \left[\frac{m^2}{z^2 f_+} - q^2 \frac{A_t^2}{h} \right] \psi = 0 \quad , \quad (24)$$

where

$$f_{\pm} = 1 + \xi \left[-3 \pm \frac{\mu^2 + \delta\mu^2}{4} z^4 \right] \quad . \quad (25)$$

The boundary behavior is altered. As $z \rightarrow 0$, we obtain $\phi \sim z^\Delta$, where

$$\Delta = \Delta_{\pm} = \frac{3}{2} \pm \sqrt{\frac{9}{4} + \frac{m^2}{1 - 3\xi}} \quad , \quad (26)$$

to be compared with (10).

We obtain modified solutions. The behavior is shown in figure 1. The figure also displays the effect of inhomogeneity (τ) on β_{max} (21) for $\xi = 0$.

The coupling to the Einstein tensor alters the near horizon limit of the theory so that

$$m^2 \rightarrow \frac{m^2 + \tau f_-(1)}{f_+(1)} \quad . \quad (27)$$

The minimum effective charge is found by setting $T = 0$. Then $\mu_{\text{eff}}^2 = \mu^2 + \delta\mu^2 = 12$, and $f_{\pm}(1) = 1 + \xi(-3 \pm 3)$. We deduce

$$q_{\text{min}}^2 = \frac{3 + 2\Delta(\Delta - 3) + 2(1 - 6\xi)\tau}{4} \quad , \quad (28)$$




FIG. 1. The critical temperature *vs.* the magnetic field numerically calculated with $q = 10$ and $\Delta = 5/2$. The dotted lines are calculated with $\xi = 0$ while the solid use $\xi = .20$. Starting from the top, on the vertical axis, the lines are $\frac{\tau}{(q\mu)^2} = 0, .05, .10, .15, .25$, and $.35$.




FIG. 2. The top line on the left-hand side of the graph corresponds to the homogeneous solution, with lines $\frac{\tau}{(q\mu)^2} = .15, .35$ below. The critical temperature of the homogeneous solution is found to decrease below the inhomogeneous lines for large β . We used $q = 10$, $\Delta = 5/2$, and $\xi = .20$.

to be compared with (22). For

$$\xi > \frac{1}{6} \quad , \quad (29)$$

the minimum charge *decreases*, and therefore the maximum value of β (21) *increases* compared to the value in the homogeneous case ($\tau = 0$). Thus, there is a neighborhood near zero temperature in which the inhomogeneous solution has higher critical temperature than the homogeneous one. Figure 2 displays the effect of the potential on the homogeneous solution as well as two inhomogeneous cases.

In conclusion, we have developed a gravitational dual theory for the FFLO state of condensed matter. The gravitational theory consists of two $U(1)$ gauge fields and a scalar coupled to a charged AdS black hole. The first gauge field produces the instability for a condensate to

form, while the second controls chemical potential associated with spin. In the absence of an interaction of the Einstein tensor with the scalar field, the system possesses dominant homogeneous solutions for all allowed values of the spin chemical potential. In the presence of the interaction term, at low temperatures, the system is shown to possess a higher critical temperature for a scalar field with spatial modulation compared to the homogeneous

solution.

ACKNOWLEDGMENTS

J. A. acknowledges support from the Office of Research at the University of Michigan-Flint. G. S. is supported by the US Department of Energy under Grant No. DE-FG05-91ER40627.

-
- [1] S. A. Hartnoll, C. P. Herzog, and G. T. Horowitz, *Phys. Rev. Lett.* **101**, 031601 (2008); S. A. Hartnoll, C. P. Herzog, and G. T. Horowitz, *JHEP* **0812**, 015 (2008).
 - [2] S. Bhattacharyya, V. E. Hubeny, S. Minwalla and M. Rangamani, *JHEP* **0802**, 045 (2008).
 - [3] M. Cubrovic, J. Zaanen and K. Schalm, *Science* **325**, 439 (2009).
 - [4] I. Martin, D. Podolsky, and S. A. Kivelson, *Phys. Rev. B* **72** (2005) 060502.
 - [5] E. Berg, E. Fradkin, S. A. Kivelson, and J. Tranquada, *New J. Phys.* **11** (2009) 115004.
 - [6] R. Flauger, E. Pajer and S. Papanikolaou, *Phys. Rev. D* **83**, 064009 (2011).
 - [7] J. A. Hutasoit, S. Ganguli, G. Siopsis and J. Therrien, *JHEP* **1202**, 086 (2012).
 - [8] S. Ganguli, J. A. Hutasoit and G. Siopsis, arXiv:1205.3107 [hep-th].
 - [9] A. Donos and J. P. Gauntlett, *JHEP* **1108**, 140 (2011).
 - [10] R. Casalbuoni and G. Nardulli, *Rev. Mod. Phys.* **76**, 263 (2004).
 - [11] P. Fulde and R. A. Ferrell, *Phys. Rev.* **135**, A550 (1964).
 - [12] A. I. Larkin and Y. N. Ovchinnikov, *Zh. Eksp. Teor. Fiz.* **47**, 1136 (1964) [*Sov. Phys. JETP* **20**, 762 (1965)].
 - [13] A. I. Buzdin, *Rev. Mod. Phys.* **77**, 935 (2005).
 - [14] T. Albash and C. V. Johnson, *JHEP* **0809**, 121 (2008).
 - [15] T. Albash and C. V. Johnson, *Phys. Rev. D* **80**, 126009 (2009); T. Albash and C. V. Johnson, arXiv:0906.0519 [hep-th].
 - [16] M. Montull, A. Pomarol and P. J. Silva, *Phys. Rev. Lett.* **103**, 091601 (2009).
 - [17] K. Maeda, M. Natsuume and T. Okamura, *Phys. Rev. D* **81**, 026002 (2010).
 - [18] X. -H. Ge, B. Wang, S. -F. Wu and G. -H. Yang, *JHEP* **1008**, 108 (2010).
 - [19] D. Arean, M. Bertolini, C. Krishnan, and T. Prochazka, *JHEP* **1109**, 131 (2011).
 - [20] F. Bigazzi, A. L. Cotrone, D. Musso, N. P. Fokeeva and D. Seminara, *JHEP* **1202**, 078 (2012).
 - [21] J. Alsup, E. Papantonopoulos and G. Siopsis, arXiv:1208.4582 [hep-th].
 - [22] A. Nicolis, R. Rattazzi, and E. Trincherini, *Phys. Rev. D* **79**, 064036 (2009);
 - [23] S. V. Sushkov, *Phys. Rev. D* **80**, 103505 (2009).
 - [24] E. N. Saridakis, S. V. Sushkov *Phys. Rev. D* **81**, 083510 (2010); C. Germani, A. Kehagias, *Phys. Rev. Lett.* **105**, 011302 (2010).
 - [25] K. Maeda, N. Ohta, and R. Wakebe, *Eur. Phys. J. C* **72** 1949 (2012).
 - [26] F. Farakos, C. Germani, A. Kehagias and E. N. Saridakis, *JHEP* **1205**, 050 (2012).
 - [27] T. Kolyvaris, G. Koutsoumbas, E. Papantonopoulos, and G. Siopsis, *Class. Quant. Grav.* **29**, 205011 (2012).
 - [28] C. Song-Bai, P. Qi-Yuan, and J. Ji-Liang, *Chin. Phys. B* **21**, 040403 (2012).
 - [29] P. Breitenlohner and D. Z. Freedman, *Annals Phys.* **144**, 249 (1982).
 - [30] G. T. Horowitz and M. M. Roberts, *JHEP* **0911**, 015 (2009).
 - [31] J. Alsup, G. Siopsis, and J. Therrien, *Phys. Rev. D* **86**, 025002 (2012).
 - [32] B. S. Chandrasekhar, *Appl. Phys. Lett.* **1**, 7 (1962). A. M. Clogston, *Phys. Rev. Lett.* **9**, 266 (1962).